




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/15, 1/21, 15/62, 15/63, 15/67, 9/42, C12P 1/02, 1/04, 1/06 // C12R 1:885		A1	(11) International Publication Number: WO 99/46362
			(43) International Publication Date: 16 September 1999 (16.09.99)
(21) International Application Number: PCT/CA99/00197		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 9 March 1999 (09.03.99)		<p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(30) Priority Data: 09/037,524 10 March 1998 (10.03.98) US			
(71) Applicant: IOGEN CORPORATION [CA/CA]; 400 Hunt Club Road, Ottawa, Ontario K1V 1C1 (CA).			
(72) Inventors: WHITE, Theresa, C.; 61 Kimberwick Crescent, Ottawa, Ontario K1V 0W6 (CA). HINDLE, Christopher, D.; 1745 Heron Road, Ottawa, Ontario K1V 6A2 (CA).			
(74) Agents: SECHLEY, Konrad, A. et al.; Gowling, Strathy & Henderson, Suite 2600, 160 Elgin Street, Ottawa, Ontario K1P 1C3 (CA).			

(54) Title: GENETIC CONSTRUCTS AND GENETICALLY MODIFIED MICROBES FOR ENHANCED PRODUCTION OF BETA-GLUCOSIDASE



(57) Abstract

This invention relates to the genetic modification of a microbe to enhance its production of an enzyme, beta-glucosidase, that is important in the cellulose conversion process. The inventors have discovered genetic constructs that, when expressed in recombinant microbes, dramatically increase the amount of beta-glucosidase produced relative to untransformed microbes. The genetic constructs comprise a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region. The increased level of beta-glucosidase significantly increases the efficiency of hydrolysis of cellulose to glucose by cellulase enzymes, thereby enhancing the production of fuel ethanol from cellulose.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**GENETIC CONSTRUCTS AND GENETICALLY MODIFIED MICROBES FOR
ENHANCED PRODUCTION OF BETA-GLUCOSIDASE**

BACKGROUND OF THE INVENTION

1. Field of the Invention

This invention relates to genetic modification of microbes to enhance production of a commercially important enzyme, beta-glucosidase. This invention also relates to genetic constructs that dramatically increase the amount of beta-glucosidase produced by microbes containing these constructs.

2. Background of the Related Art

The possibility of producing ethanol from cellulose has received much attention due to the availability of large amounts of feedstock, the desirability of avoiding burning or land filling the materials, and the cleanliness of the ethanol fuel. The advantages of such a process for society are described in the cover story of the Atlantic Monthly, April 1996.

The natural cellulosic feedstocks for such a process are referred to as "biomass". Many types of biomass, including wood, agricultural residues, herbaceous crops, and municipal solid wastes, have been considered as feedstocks for ethanol production. These materials primarily consist of cellulose, hemicellulose, and lignin. This invention can be applied to the conversion of the cellulose to ethanol.

Cellulose is a polymer of the simple sugar glucose connected by beta 1,4 linkages. Cellulose is very resistant to degradation or depolymerization by acid, enzymes, or micro-organisms. Once the cellulose is converted to glucose, the resulting sugar is easily fermented to ethanol using yeast. The difficult challenge of the process is to convert the cellulose to glucose.

The oldest methods studied to convert cellulose to glucose are based on acid hydrolysis (review by Grethlein, "Chemical breakdown of Cellulosic Materials", J. Appl. Chem. Biotechnol. 28:296-308 (1978)). This process can involve the use of concentrated or dilute acids. The concentrated acid process produces a high yield of glucose, but the recovery of the acid, the specialized materials of construction required, and the need to minimize water in the system are serious disadvantages of this process. The dilute acid process uses low levels of acid to overcome the need for chemical recovery. However, the maximum glucose yield is only about 55% of the cellulose, and a high degree of production of degradation products can inhibit the fermentation to ethanol by yeast. These problems have prevented the acid hydrolysis process from reaching commercialization.

To overcome the problems of the acid hydrolysis process, cellulose conversion processes have focused more recently on enzymatic hydrolysis, using cellulase enzymes. Enzymatic hydrolysis of cellulose is carried out by mixing the substrate and water to achieve a slurry of 5% to 12% cellulose and adding 5 to 50 International Units (IU) cellulase enzymes per gm cellulose. Typically, the hydrolysis is run for 12 to 150 hours at 35-60°C, pH 4-6.

Many microbes make enzymes that hydrolyze cellulose, including the wood rotting fungus *Trichoderma*, the compost bacteria *Thermomonospora*, *Bacillus*, and *Cellulomonas*; *Streptomyces*; and the fungi *Humicola*, *Aspergillus* and *Fusarium*. The enzymes made by these microbes are mixtures of proteins with three types of actions useful in the conversion of cellulose to glucose: endoglucanases (EG), cellobiohydrolases (CBH), and beta-glucosidase. EG and CBH enzymes are collectively referred to as "cellulase."

EG enzymes cut the cellulose polymer at random locations, opening it up to attack by CBH enzymes. As an example, *Trichoderma* strains produce at least four distinct EG enzymes, known as EGI, EGII, EGIII, and EGV.

CBH enzymes sequentially release molecules of cellobiose from the ends of the cellulose polymer. Cellobiose is the water-soluble beta-1,4-linked dimer of glucose. There are two primary CBH enzymes made by *Trichoderma*, CBHI and CBHII.

Beta-glucosidase enzymes hydrolyze cellobiose to glucose. *Trichoderma* makes one beta-glucosidase enzyme.

This final step in the cellulose hydrolysis which is catalyzed by beta-glucosidase is important, because glucose is readily fermented to ethanol by a variety of yeasts while cellobiose is not. Any cellobiose remaining at the end of the hydrolysis represents a loss of yield of ethanol. More importantly, cellobiose is an extremely potent inhibitor of the CBH and EG enzymes. Cellobiose decreases the rate of hydrolysis of the *Trichoderma* CBH and EG enzymes by 50% at a concentration of only 3.3 g/L. The decrease in rate of hydrolysis necessitates the addition of higher levels of cellulase enzymes, which adversely impacts the overall process economics. Therefore, the accumulation of cellobiose during hydrolysis is extremely undesirable for ethanol production.

Cellobiose accumulation has been a major problem in enzymatic hydrolysis because *Trichoderma* and the other cellulase-producing microbes make very little beta-glucosidase. Less than 1% of the total protein made by *Trichoderma* is beta-glucosidase. The low amount of beta-glucosidase results in a shortage of capacity to hydrolyze the cellobiose to glucose and an accumulation of 10 to 20 g/L of cellobiose during hydrolysis. This high level of cellobiose increases the amount of cellulase required by 10-fold over that if an adequate amount of beta-glucosidase were present.

Several approaches have been proposed to overcome the shortage of beta-glucosidase in cellulase enzymes.

One approach has been to produce beta-glucosidase using microbes that produce little cellulase, and add this beta-glucosidase exogenously to cellulase enzyme to enhance the hydrolysis. The most successful of such beta-glucosidase producing microbes have been *Aspergillus niger* and *Aspergillus phoenicis*. Beta-glucosidase from these microbes are available commercially as Novozym 188 from Novo Nordisk. However, the quantities required are much too costly for a commercial biomass to ethanol operation.

A second approach to overcoming the shortage of beta-glucosidase is to carry out cellulose hydrolysis simultaneously with fermentation of the glucose by yeast, the so-called simultaneous saccharification and fermentation (SSF) process. In an SSF system, the fermentation of the glucose removes it from solution. Glucose is a potent inhibitor of beta-glucosidase, so SSF is an attempt to increase the efficiency of beta-glucosidase. However, SSF systems are not yet commercially viable because the operating temperature for yeast of 28°C is too low for the 50°C conditions required by cellulase; operation at a compromise temperature of 37°C is inefficient and prone to microbial contamination.

A third approach to overcoming the shortage of beta-glucosidase is to use genetic engineering to overexpress the enzyme and increase its production by *Trichoderma*. This approach was taken by Barnett, Berka, and Fowler, in "Cloning and Amplification of the Gene Encoding an Extracellular β -glucosidase from *Trichoderma reesei*: Evidence for Improved Rates of Saccharification of Cellulosic Substrates," Bio/Technology, Volume 9, June 1991, p. 562-567, herein referred to as "Barnett, et al."; and Fowler, Barnett, and Shoemaker in WO 92/10581, "Improved Saccharification of Cellulose by Cloning and Amplification

of the β -glucosidase gene of *Trichoderma reesei*," herein referred to as "Fowler, et al."

Both Barnett, et al. and Fowler, et al. describe the insertion of multiple copies of the beta-glucosidase gene into *Trichoderma reesei* strain P40. Both groups constructed plasmid pSAS β -glu, a transformation vector containing the genomic *T. reesei* beta-glucosidase gene and the *amdS* selectable marker. The *amdS* gene is from *Aspergillus nidulans* and codes for the enzyme acetamidase, which allows transformed cells to grow on acetamide as a sole source of nitrogen. *T. reesei* does not contain a functional equivalent to the *amdS* gene and is therefore unable to utilize acetamide as a nitrogen source. The transformed cells contained 10 to 15 copies of the beta-glucosidase gene and produced 5.5-fold more beta-glucosidase than the untransformed cells.

The enhanced production of beta-glucosidase obtained by Barnett, et al. and Fowler, et al. is not sufficient to alleviate the shortage of beta-glucosidase for cellulose hydrolysis. The amount of beta-glucosidase made by natural *Trichoderma* strains, for example, must be increased at least 10-fold to meet the requirements of cellulose hydrolysis.

When overexpressing proteins in *Trichoderma*, one strategy is to link the gene of interest directly to the *cbh1* promoter or to the *cbh1* secretion signal. Since CBH1 is the most abundant protein produced by *Trichoderma* under cellulase-inducing conditions, the *cbh1* promoter and secretion signal are thought to be very effective in directing the transcription and secretion of proteins encoded by a gene positioned after them in a genetic construct. Such a strategy has been successfully used to overexpress proteins from *Trichoderma* and other microorganisms (Margolles-Clark, Hayes, Harman and Penttila, 1996, "Improved Production of *Trichoderma harzianum* endochitinase by expression in *Trichoderma reesei*", Appl. Environ. Microbiol. 62(6): 2145-2151; Joutsjouki, Torkkeli and Nevalainen, 1993, "Transformation of *Trichoderma reesei* with the

Hormoconis resinae glucoamylase P (*gamP*) gene: production of a heterologous glucoamylase by *Trichoderma reesei*", Curr. Genet. 24: 223-228; Karhunen, Mantyla, Nevalainen and Suominen, 1993, "High frequency one-step gene replacement in *Trichoderma reesei* 1. Endoglucanase I overproduction", Mol. Gen. Genet. 241: 515-522).

In spite of a significant amount of research effort, there has not been a means to produce sufficiently high levels of beta-glucosidase. Such a process would be a large step forward in the production of fuel alcohol from cellulose.

SUMMARY OF THE INVENTION

The inventors have made a discovery that enables the production of beta-glucosidase enzyme at levels far higher than currently achievable. The high levels of beta-glucosidase improve the efficiency of the enzymatic hydrolysis of cellulose to glucose. The resulting decrease in enzyme requirement, increased cellulose conversion, decrease in hydrolysis time, or a combination of these advantages, decreases the overall process costs of converting cellulose to ethanol.

The inventors have discovered genetic constructs that significantly increase the production of beta-glucosidase by recombinant microbes in which the constructs are expressed. The genetic constructs that accomplish this task comprise DNA sequences that encode a mature beta-glucosidase enzyme and a xylanase secretion signal.

As far as the inventors are aware, there have been no previous reports that linking the xylanase secretion signal to the mature beta-glucosidase enzyme increases the production of beta-glucosidase. In addition, the inventors are not aware of previous reports that linking the xylanase secretion signal to any non-xylanase mature protein increases the production of the protein. The inventors have discovered this surprising and unreported result. It was further surprising

that the use of the xylanase secretion signal resulted in higher levels of beta-glucosidase than the use of the *cbh1* secretion signal. Since xylanase comprises a much smaller proportion of the total protein produced by *Trichoderma* than does CBH1 (5% and 60%, respectively), one would expect that the *cbh1* secretion signal would be more effective. The reasons why linking the xylanase secretion signal to the mature beta-glucosidase enzyme increases beta-glucosidase production are not known, but might relate to the similarity in length between the beta-glucosidase and xylanase secretion signals or to the lower abundance of xylanase against which the recombinant beta-glucosidase must compete for secretion out of the cell. However, the practice of the invention is not limited by these or any other specific reasons.

The present invention is not anticipated by Barnett, et al. and Fowler, et al., who each disclosed the enhanced expression of beta-glucosidase by recombinant means. The genetic construct of Barnett, et al. and Fowler, et al. comprises the beta-glucosidase promoter, coding region and secretion signal. The methods used by Barnett, et al. and Fowler, et al. are not as effective as the methods taught by the inventors, and do not anticipate the genetic constructs of the present invention.

In one aspect of our invention, a genetically modified microbe for producing beta-glucosidase comprises a beta-glucosidase construct not present in an untransformed microbe from which said genetically modified microbe is derived, said beta-glucosidase construct having a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, wherein said genetically modified microbe is selected from the group consisting of *Trichoderma*, *Humicola*, *Fusarium*, *Streptomyces*, *Thermomonospora*, *Bacillus*, *Cellulomonas*, and *Aspergillus*, and wherein said genetically modified microbe produces at least about a 10-fold increase in production of beta-glucosidase relative to said untransformed microbe.

In another aspect, our invention includes a beta-glucosidase genetic construct comprising a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, wherein said beta-glucosidase genetic construct, when introduced into and expressed in an untransformed microbial host selected from the group consisting of *Trichoderma*, *Humicola*, *Fusarium*, *Streptomyces*, *Thermomonospora*, *Bacillus*, *Cellulomonas*, and *Aspergillus*, produces at least about a 10-fold increase in production of beta-glucosidase relative to said untransformed microbial host.

In still another aspect of our invention, a genetically modified *Trichoderma* microbe for producing beta-glucosidase comprises a beta-glucosidase construct not present in an untransformed *Trichoderma* microbe, said beta-glucosidase construct having a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, wherein said genetically modified *Trichoderma* produces at least about a 10-fold increase in production of beta-glucosidase relative to said untransformed *Trichoderma* microbe.

In yet another aspect, our invention includes a genetically modified *Trichoderma reesei* microbe for producing beta-glucosidase comprising a beta-glucosidase construct not present in an untransformed *Trichoderma reesei* microbe, said beta-glucosidase construct having a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, wherein said genetically modified *Trichoderma reesei* microbe produces at least about a 10-fold increase in production of beta-glucosidase relative to said untransformed microbe.

In yet another aspect of our invention, a beta-glucosidase genetic construct comprises a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, wherein said beta-glucosidase genetic construct, when introduced into and expressed in a *Trichoderma* microbe, produces at least about a 10-fold

increase in production of beta-glucosidase relative to an untransformed *Trichoderma* microbe.

In still yet another aspect of our invention, a beta-glucosidase genetic construct comprises a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, wherein said beta-glucosidase genetic construct, when introduced into and expressed in a *Trichoderma reesei* microbe, produces at least about a 10-fold increase in production of beta-glucosidase relative to an untransformed *Trichoderma reesei* microbe.

Other aspects of our invention will be better understood and advantages thereof more apparent in view of the following detailed description of the preferred embodiments and accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Restriction map of the vector pCBG1-TV and the amino acid sequence of the CBH1 secretion signal/mature beta-glucosidase juncture.

Figure 2: Restriction map of the vector pXBG1-TV and the amino acid sequence of the xylanase II secretion signal/mature beta-glucosidase juncture.

Figure 3: Restriction map of the vector pC/XBG(Xba1)-TV and the amino acid sequence of the xylanase II secretion signal/mature beta-glucosidase juncture.

Figure 4: Southern blot of genomic DNA isolated from *T. reesei* strains RutC30 and M2C38 and probed with a labeled DNA fragment comprising the M2C38 xylanase promoter plus secretion signal.

Figure 5: Southern blot of genomic DNA isolated from *T. reesei* strains RutC30 and M2C38 and probed with a labeled DNA fragment comprising the M2C38 mature beta-glucosidase coding region.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Preferred embodiments of this invention are described by first defining the following terms.

Beta-glucosidase is an enzyme that hydrolyzes the glucose dimer cellobiose to glucose. There are many microbes that make beta-glucosidase, and the properties of these enzymes vary, including structure (molecular weight, three-dimensional orientation, amino acid composition, and active site) and catalytic activity (rate and kinetics of cellobiose hydrolysis, and ability to act on other substrates). However, in all cases beta-glucosidase enzyme can hydrolyze cellobiose to glucose. This may also be referred to as a mature beta-glucosidase enzyme when the active enzyme does not contain a beta-glucosidase secretion signal peptide.

The preferred beta-glucosidase for practicing the invention is the beta-glucosidase made by *Trichoderma*. This beta-glucosidase enzyme is of molecular weight 74,000 (as measured by SDS-polyacrylamide gel electrophoresis) and has an isoelectric point of 8.3 (as measured by non-denaturing isoelectric focusing polyacrylamide gel electrophoresis).

Beta-glucosidase gene is a region of DNA that codes for the production of beta-glucosidase enzyme. All microbes that produce beta-glucosidase possess at least one beta-glucosidase gene. A natural beta-glucosidase gene comprises a beta-glucosidase promoter, a secretion signal, a coding region and a transcriptional

terminator. Microbes that do not produce beta-glucosidase do not generally contain an active or functional beta-glucosidase gene.

Beta-glucosidase secretion signal is the DNA sequence that encodes the beta-glucosidase secretion signal peptide.

Beta-glucosidase secretion signal peptide is the peptide sequence present at the amino terminus of the encoded beta-glucosidase enzyme that is subsequently removed during export of the mature beta-glucosidase enzyme out of the microbial cells. The secretion signal may comprise a pro-peptide, a pre-peptide or both.

Mature beta-glucosidase coding region comprises the DNA sequence necessary to encode the functional beta-glucosidase enzyme, as isolated from extracellular culture filtrates, but not the beta-glucosidase secretion signal.

Xylanase is an enzyme that hydrolyzes xylan to xylose. There are many microbes that make xylanase, and the properties of these enzymes vary, including structure (molecular weight, three-dimensional orientation, amino acid composition, and active site) and catalytic activity (rate and kinetics of xylan hydrolysis, and ability to act on other substrates). However, in all cases xylanase enzyme can hydrolyze xylan to xylose. This may also be referred to as a mature xylanase enzyme when the active enzyme does not contain a xylanase secretion signal peptide.

Some of the most important commercial xylanases are classified as Family 11 xylanases. A xylanase enzyme is classified in Family 11 if it possesses the amino acids common to Family 11, including two glutamic acid residues serving as the essential catalytic residues. These residues are amino acids 86 and 177 by *Trichoderma reesei* xylanase II numbering. The amino acids common to Family 11 xylanases are described in Wakarchuck, et al, Protein Science 3:467-475 (1994).

Xylanase gene is a region of DNA that codes for the production of xylanase enzyme. All microbes that produce xylanase possess at least one xylanase gene. A natural xylanase gene comprises a xylanase promoter, a secretion signal, a coding region and a transcriptional terminator. Microbes that do not produce xylanase do not generally contain an active or functional xylanase gene.

Xylanase secretion signal is the DNA sequence that encodes the xylanase secretion signal peptide.

Xylanase secretion signal peptide is the peptide sequence present at the amino terminus of the encoded xylanase enzyme that is subsequently removed during export of the mature xylanase enzyme out of the microbial cells. The secretion signal may comprise a pro-peptide, a pre-peptide or both.

Cellulase is an enzyme that hydrolyzes cellulose to short beta 1,4-linked oligomers of glucose, including cellotetraose, cellotriose and cellobiose. There are many microbes that make one or more cellulase enzymes often classified as cellobiohydrolases or endoglucanases. The properties of these enzymes vary, including structure (molecular weight, three-dimensional orientation, amino acid composition, and active site) and catalytic activity (rate and kinetics of xylan hydrolysis, and ability to act on other substrates). However, in all cases cellulase enzymes can hydrolyze cellulose to short beta 1,4-linked oligomers of glucose, including cellotetraose, cellotriose and cellobiose.

Beta-glucosidase genetic construct refers to a gene comprising the elements necessary to produce beta-glucosidase. These include:

- a. A mature beta-glucosidase coding region.

In a preferred embodiment, the mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* gene. The DNA sequence of the mature beta-glucosidase coding region from *Trichoderma reesei* can be found in Figure 1 of Barnett, et al.

Those skilled in the art are aware that a natural structural region can be modified by replacement, substitution, addition, or elimination of one or more nucleic acids without changing its function. The practice of the invention encompasses and is not constrained by such alterations to the mature beta-glucosidase coding region.

b. A xylanase secretion signal.

In a preferred embodiment, the xylanase secretion signal comprises a xylanase secretion signal of a Family 11 xylanase gene.

In a more preferred embodiment, the Family 11 xylanase gene is a *Trichoderma* xylanase gene.

In an even more preferred embodiment, the xylanase secretion signal comprises a xylanase secretion signal of *Trichoderma* xylanase I (*xln1*) gene or xylanase II (*xln2*) gene. The DNA sequences of the *Trichoderma xln1* and *xln2* secretion signals can be found in Figures 3 and 2, respectively, of Torronen, Mach, Messner, Gonzalez, Kalkkinen, Harkki and Kubicek, "The two major xylanases from *Trichoderma reesei*: characterization of both enzymes and genes," Bio/Technology 10: 1461-1465, 1992 (the gene identifications in the figure legends, as published, are reversed).

Those skilled in the art are aware that a natural secretion signal can be modified by replacement, substitution, addition, or elimination of one or more nucleic

acids without changing its function. The practice of the invention encompasses and is not constrained by such alterations to the xylanase secretion signal.

c. Promoter

The practice of the invention is not constrained by the choice of promoter in the genetic construct. However, preferred promoters are the *Trichoderma cbh1*, *cbh2*, *eg1*, *eg2*, *eg3*, *eg5*, *xln1* and *xln2* promoters. The DNA sequence of the *Trichoderma reesei cbh1* is deposited in GenBank under Accession Number D86235.

Those skilled in the art are aware that a natural promoter can be modified by replacement, substitution, addition or elimination of one or more nucleotides without changing its function. The practice of the invention encompasses and is not constrained by such alterations to the promoter.

d. Additional sequences between the xylanase secretion signal and the mature beta-glucosidase coding region

The genetic constructs described in Examples 5, 6 and 7 contain nine additional base pairs of DNA sequence as shown in Figures 1-3; the first three encode the glutamine residue after the secretion signal of the *Trichoderma reesei* xylanase II gene, and the remaining six result from the insertion and/or modification of unique restriction sites used to join the xylanase secretion signal to the mature beta-glucosidase coding region. These DNA sequences result in the presence of additional amino acids between the xylanase secretion signal peptide and the mature beta-glucosidase enzyme. These DNA sequences, which may be natural or synthetic, may encode one or more of the amino acids of the mature xylanase protein corresponding to the xylanase secretion signal encoded by the construct or may result from the addition of restriction enzyme sites needed to join the xylanase secretion signal peptide and mature beta-glucosidase enzyme. The

practice of the invention encompasses but is not constrained by the presence of additional DNA sequences between the xylanase secretion signal and the mature beta-glucosidase coding region.

e. other elements

The genetic construct contains a transcriptional terminator immediately downstream of the mature beta-glucosidase coding region. The practice of the invention is not constrained by choice of transcriptional terminator and may include as much DNA downstream (i.e., at the 3' end) of the stop codon of any known coding region as is sufficient to direct the termination of transcription by RNA polymerase. The transcriptional terminator present downstream of the mature beta-glucosidase coding region in the constructs described in Examples 5-7 comprises 1.9 kb of DNA 3' to the stop codon of the *Trichoderma cbh2* gene.

The DNA sequence of the first 553 base pairs of the *Trichoderma reesei cbh2* transcriptional terminator, which are located immediately downstream (or 3') of the TAA stop codon, can be found in Figure 2 of Chen, Gritzali and Stafford, "Nucleotide Sequence and Deduced Primary Structure of Cellobiohydrolase II from *Trichoderma reesei*," Bio/Technology 5: 274-278, 1987.

The genetic construct contains a selectable marker which may be present upstream or downstream of the genetic construct (i.e., at the 5' or 3' end of the construct) on the same plasmid vector or may be cotransformed with the construct on a separate plasmid vector. Choices of selectable markers are well known to those skilled in the art and include genes (synthetic or natural) that confer to the transformed cells the ability to utilize a metabolite that is not normally metabolized by the microbe (e.g., the *A. nidulans amdS* gene encoding acetamidase and conferring the ability to grow on acetamide as the sole nitrogen source) or antibiotic resistance (e.g., the *Escherichia coli hph* gene encoding hygromycin-b-phosphotransferase and conferring resistance to hygromycin). If

the host strain lacks a functional gene for the marker chosen, then that gene may be used as a marker. Examples of such markers include *trp*, *pyr4*, *pyrG*, *argB*, *leu*, and the like. The corresponding host strain would therefore have to lack a functional gene corresponding to the marker chosen, i.e., *trp*, *pyr*, *arg*, *leu* and the like. The selectable marker used in the genetic constructs described in Examples 5-7 is the *E. coli hph* gene expressed from the *Trichoderma* phosphoglycerate kinase (*pgk*) promoter.

The DNA sequence of the *E. coli hph* gene can be found in Figure 4 of Gritz and Davies, "Plasmid-encoded hygromycin B resistance: the sequence of hygromycin B phosphotransferase gene and its expression in *Escherichia coli* and *Saccharomyces cerevisiae*," Gene 25: 179-188, 1983; the DNA sequence of the *Trichoderma reesei pgk* promoter can be found in Figure 2 of Vanhanen, Saloheimo, Ilmen, Knowles and Penttila, "Promoter structure and expression of the 3-phosphoglycerate kinase-encoding gene (*pgk1*) of *Trichoderma reesei*," Gene 106: 129-133, 1991.

One preferred embodiment of the invention comprises the beta-glucosidase genetic construct described thus far. The practice of our invention is not constrained by the method of making the construct, which can include, but is not restricted to, standard molecular biology techniques such as isolation of plasmid DNA from *E. coli* by alkaline lysis, digestion of plasmid DNA with restriction endonucleases, separation and isolation of DNA fragments by agarose gel electrophoresis, ligation of DNA fragments with T4 DNA ligase, insertion of unique restriction sites at the ends of DNA fragments by polymerase chain reaction or the addition of oligonucleotide linkers, and the blunting of DNA fragments with T4 DNA polymerase or Klenow fragment of *E. coli* DNA polymerase I.

Examples 1-7 describe procedures for making such genetic constructs.

In another preferred embodiment of our invention, the beta-glucosidase genetic construct is introduced into and expressed in a microbial host to create a genetically modified microbe. The resulting genetically modified microbe produces an increased level of beta-glucosidase relative to the untransformed microbial host. The genetically modified microbe produces an increased level of beta-glucosidase of preferably at least about 10-fold relative to the untransformed microbial host, more preferably at least about 40-fold relative to the untransformed microbial host, and most preferably at least about 120-fold relative to the untransformed microbial host.

This invention encompasses any method of introducing the beta-glucosidase genetic construct into the microbial host familiar to those skilled in the art, including but not limited to, calcium chloride treatment of bacterial cells or fungal protoplasts to weaken the cell membranes, addition of polyethylene glycol to allow for fusion of cell membranes, depolarization of cell membranes by electroporation, or shooting the DNA through the cell wall and membranes via microprojectile bombardment with a particle gun.

Example 8 describes the procedures for introducing the beta-glucosidase genetic construct into *Trichoderma* spores using a particle gun.

A 10-fold enhancement of beta-glucosidase production relative to the untransformed microbial host reflects a significant enhancement that is well above the natural variability of the strain and commercially significant. The degree of enhancement of beta-glucosidase by this method has been as high as 126-fold and could reach over 1000-fold. The measurement of the degree of enhancement of beta-glucosidase production is by growth of the culture and measurement of the beta-glucosidase activity, as described in Example 11. It is believed that genetic constructs of our invention will produce any level of enhancement greater than about 10-fold.

It is understood by those skilled in the art that the specific beta-glucosidase activity of an enzyme mixture (in IU/mg protein) can be increased by decreasing the amount of cellulase and other proteins in the enzyme mixture. This can be done by physical and mechanical separations of the enzyme mixture or by deletion of the cellulase or other genes by recombinant means. Such methods have little or no effect on the actual production of beta-glucosidase by the microorganism. These procedures may, however, be optionally included in the practice of our invention.

In a preferred embodiment, the microbial host is a member of the species of *Trichoderma*, *Humicola*, *Fusarium*, *Aspergillus*, *Streptomyces*, *Thermomonospora*, *Bacillus*, or *Cellulomonas*. These species are well suited because they produce cellulase in addition to beta-glucosidase. In addition, methods have been published for the introduction of DNA constructs into cellulase-producing strains of *Trichoderma* (Lorito, Hayes, DiPietro and Harman, 1993, "Biolistic Transformation of *Trichoderma harzianum* and *Gliocladium virens* using plasmid and genomic DNA," Curr. Genet. 24: 349-356; Goldman, VanMontagu and Herrera-Estrella, 1990, "Transformation of *Trichoderma harzianum* by high-voltage electric pulse", Curr. Genet. 17:169-174; Penttila, Nevalainen, Ratto, Salminen and Knowles, 1987, "A versatile transformation system for the cellulolytic fungus *Trichoderma reesei*", Gene 6: 155-164), *Aspergillus* (Yelton, Hamer and Timberlake, 1984, "Transformation of *Aspergillus nidulans* using a *trpC* plasmid," Proc. Natl. Acad. Sci. USA 81: 1470-1474), *Fusarium* (Bajar, Podila and Kolattukudy, 1991, "Identification of a fungal cutinase promoter that is inducible by a plant signal via a phosphorylated trans-acting factor," Proc. Natl. Acad. Sci. USA 88: 8202-8212), *Streptomyces* (Hopwood et al., 1985, "Genetic Manipulation of *Streptomyces*: a laboratory manual," The John Innes Foundation, Norwich, UK) and *Bacillus* (Brigidi, DeRossi, Bertarini, Riccardi and Matteuzzi, 1990, "Genetic transformation of intact cells of *Bacillus subtilis* by electroporation," FEMS Microbiol. Lett. 55: 135-138).

The genetic constructs used in these published transformation methods are similar to those described in Examples 5-7 in that they contain a promoter linked to a protein coding region (which may encode a selectable marker) and a transcriptional terminator. In most cases, the genetic constructs are linked to a selectable marker gene.

Although there are no published methods for the transformation of *Humicola*, *Thermomonospora* or *Cellulomonas*, it is believed that the transformation methods for other filamentous fungi or bacteria can be optimized for *Humicola*, *Thermomonospora* or *Cellulomonas* strains by virtue of the similar morphologies and physiologies of these species to those for which transformation methods have been published. In addition, transformation methods such as electroporation and particle bombardment have been used to introduce DNA into many different cell types including mammalian and plant cells, bacterial, yeast, and fungal cells.

In a preferred embodiment, the xylanase secretion signal is native to the microbial host from which said genetically modified microbe is derived (i.e., the source of the xylanase secretion signal is the same type of microbial host as the microbial host from which said genetically modified microbe is derived).

In a more preferred embodiment, the microbial host is *Trichoderma*.

In a more preferred embodiment, the microbial host is *Trichoderma reesei*.

EXAMPLES

Example 1 describes the isolation of genomic DNA from *Trichoderma reesei* strains RutC30, M2C38, BTR48 and the genetically modified derivatives of these

strains. Examples 2-7 describe the construction of genomic DNA libraries, the cloning of various genes, and several genetic constructs from *Trichoderma reesei* strain M2C38. Examples 9 and 11-15 describe the transformation and expression of beta-glucosidase genetic constructs in *Trichoderma reesei* strains M2C38, BTR48, and RutC30.

Trichoderma reesei strains M2C38 and BTR48 are proprietary strains of Iogen Corporation, and were derived from *Trichoderma reesei* RutC30 (ATCC 56765, Montenecourt and Eveleigh, 1979, "Selective isolation of high yielding cellulase mutants of *T. reesei*", Adv. Chem. Ser. 181: 289-301), which was in turn derived from *Trichoderma reesei* Qm6A (ATCC 13631 Mandels and Reese, 1957 "Induction of cellulase in *Trichoderma viride* as influenced by carbon sources and metals", J. Bacteriol. 73: 269-278).

In Example 1 and subsequent Examples, restriction endonucleases, T4 DNA polymerase, T4 DNA ligase and Klenow fragment of *E. coli* DNA polymerase 1 were purchased from Gibco/BRL, New England Biolabs, Boehringer Mannheim or Pharmacia and used as recommended by the manufacturer. *Pwo* polymerase with proof-reading activity (Boehringer Mannheim) was used in all polymerase-chain reactions (PCR) according to the manufacturer's protocol. Hygromycin B was purchased from CalBiochem.

EXAMPLE 1

Isolation of *Trichoderma reesei* Genomic DNA

To isolate genomic DNA, 50 ml of Potato Dextrose Broth (Difco) was inoculated with *T. reesei* spores collected from a Potato Dextrose Agar plate with a sterile inoculation loop. The cultures were shaken at 200 rpm for 2-3 days at 28°C. The mycelia was filtered onto a sterile GFA glass microfibre filter (Whatman) and washed with cold, deionized water. The fungal cakes were frozen in liquid

nitrogen and crushed into a powder with a pre-chilled mortar and pestle; 0.5 g of powdered biomass were resuspended in 5 ml of 100 mM Tris, 50 mM EDTA, pH 7.5 plus 1% sodium dodecyl sulphate (SDS). The lysate was centrifuged (5000g for 20 min, 4°C) to pellet cell debris. The supernatant was extracted with 1 volume buffer (10 mM Tris, 1 mM EDTA, pH 8.0) saturated phenol followed by extraction with 1 volume of buffer-saturated phenol:chloroform:isoamyl alcohol (25:24:1) in order to remove soluble proteins. DNA was precipitated from the solution by adding 0.1 volume of 3M sodium acetate, pH 5.2 and 2.5 volumes of cold 95% ethanol. After incubating for at least 1 h at -20°C, the DNA was pelleted by centrifugation (5000g for 20 min, 4°C), rinsed with 10 ml 70% ethanol, air-dried and resuspended in 1 ml 10mM Tris, 1 mM EDTA, pH8.0. RNA is digested by the addition of Ribonuclease A (Boehringer Mannheim) added to a final concentration of 0.1 mg/ml and incubation at 37°C for 1 hour. Sequential extractions with 1 volume of buffer-saturated phenol and 1 volume of buffer-saturated phenol:chloroform:isoamyl alcohol (25:24:1) are used to remove the ribonuclease from the DNA solution. The DNA is again precipitated with 0.1 volume of 3M sodium acetate, pH 5.2 and 2.5 volumes of cold 95% ethanol, pelleted by centrifugation, rinsed with 70% ethanol, air-dried and resuspended in 50 µl of 10 mM Tris, 1 mM EDTA, pH 8.0. The concentration of DNA was determined by measuring the absorbance of the solution at 260nm (p. C1 in Sambrook, Fritsch and Maniatis, "Molecular Cloning: A Laboratory Manual, Second Edition", Cold Spring Harbor Press 1989, hereafter referred to as Sambrook et al.).

EXAMPLE 2

Construction of *T. reesei* Genomic Libraries

Two plasmid libraries and one phage library were constructed using genomic DNA isolated from *T. reesei* strain M2C38. The plasmid libraries were constructed in the vector pUC119 (Viera and Messing, "Isolation of single-stranded plasmid DNA", Methods Enzymol. 153:3, 1987) as follows: 10 µg genomic DNA was digested for 20 hrs at 37°C in a 100 µl volume with 2 units/µg of *Hind*III, *Bam*HI or *Eco*RI restriction enzymes. The digested DNA was fractionated on a 0.75% agarose gel run in 0.04M Tris-acetate, 1 mM EDTA, and stained with ethidium bromide. Gel slices corresponding to the sizes of the genes of interest (based on published information and Southern blots) were excised and subjected to electro-elution to recover the DNA fragments (Sambrook et al., pp. 6.28-6.29). These enriched fractions of DNA were ligated into pUC119 in order to create gene libraries in ligation reactions containing 20-50 µg/ml DNA in a 2:1 molar ratio of vector:insert DNA, 1 mM ATP and 5 units T4 DNA ligase in a total volume of 10-15 µl at 4°C for 16 h. *Escherichia coli* strain HB101 was electroporated with the ligation reactions using the Cell Porator System (Gibco/BRL) following the manufacturer's protocol and transformants selected on LB agar containing 70 µg/ml ampicillin.

The phage library was constructed in the lambda vector lambda DASH (Stratagene, Inc.) as follows: genomic DNA (3 µg) was digested with 2, 1, 0.5 and 0.2 units/µg *Bam* HI for 1 hour at 37°C to generate fragments 9-23 kB in size. The DNA from each digest was purified by extraction with 1 volume Tris-saturated phenol:chloroform:isoamyl alcohol (25:24:1) followed by precipitation with 10 µl 3M sodium acetate, pH 5.2 and 250 µl 95% ethanol (-20°C). The digested DNA was pelleted by microcentrifugation, rinsed with 0.5 ml cold 70% ethanol, air-dried and resuspended in 10 µl sterile, deionized water. Enrichment of DNA fragments 9-23 kB in size was confirmed by agarose gel electrophoresis (0.8% agarose in 0.04 M Tris-acetate, 1 mM EDTA). Digested DNA (0.4 µg) was ligated

to 1 µg lambdaDASH arms predigested with *Bam*HI (Stratagene) in a reaction containing 2 units T4 DNA ligase and 1 mM ATP in a total volume of 5 µl at 4°C overnight. The ligation mix was packaged into phage particles using the GigaPack® II Gold packaging extracts (Stratagene) following the manufacturer's protocol. The library was titered using the *E. coli* host strain XL1-Blue MRA (P2) and found to contain 3×10^5 independent clones.

EXAMPLE 3

Isolation of *T. reesei* M2C38 clones of the cellobiohydrolase I (*cbh1*), cellobiohydrolase II (*cbh2*) and β-glucosidase (*bgl1*) genes from the pUC119 libraries

E. coli HB101 transformants harboring *cbh1*, *cbh2* or *bgl1* clones from the recombinant pUC119-*Hind* III, -*Bam*HI or -*Eco*RI libraries were identified by colony lift hybridization: $1-3 \times 10^4$ colonies were transferred onto HyBond™ nylon membranes (Amersham); membranes were placed colony-side up onto blotting paper (VWR 238) saturated with 0.5 M NaOH, 1 M NaCl for 5 min to lyse the bacterial cells and denature the DNA; the membranes were then neutralized by placing them colony-side up onto blotting paper (VWR 238) saturated with 1.5 M Tris, pH 7.5 plus 1 M NaCl for 5 min; the membranes were allowed to air-dry for 30 min and the DNA was then fixed to the membranes by baking at 80°C for 2 h.

³²P-labelled probes were prepared by PCR amplification of short (0.7-1.5 kB) fragments of the *bgl1*, *cbh1* and *cbh2* coding regions from the enriched pool of *Hind* III, *Bam*HI or *Eco*RI fragments, respectively, in a labeling reaction containing 10-50 ng target DNA, 0.2 mM each d(GCT)TP, 0.5 µM dATP, 20-40 µCi alpha-³²P-dATP, 10 pmole oligonucleotide primers and 0.5 units Taq polymerase in a total volume of 20 µl. The reaction was subjected to 6-7 cycles of amplification (95°C, 2 min; 56°C, 1.5 min; 70°C, 5 min). The amplified, ³²P-labelled DNA was precipitated by the addition of 0.5 ml 10% (w/v) trichloroacetic

acid and 0.5 mg yeast tRNA. The DNA was pelleted by microcentrifugation, washed twice with 1 ml 70% ethanol, air-dried and resuspended in 1M Tris pH7.5, 1mM EDTA.

Nylon membranes onto which the recombinant pUC119 plasmids had been fixed were prehybridized in heat-sealed bags for 1 h at 60-65°C in 1 M NaCl, 1% SDS, 50 mM Tris, 1mM EDTA pH 7.5 with 100 µg/ml denatured sheared salmon sperm DNA. Hybridizations were performed in heat-sealed bags in the same buffer with only 50 µg/ml denatured sheared salmon sperm DNA and 5×10^6 - 5×10^7 cpm of denatured *bgl1*, *cbh1* or *cbh2* probe for 16-20 h at 60-65°C. Membranes were washed once for 15 min with 1 M NaCl, 0.5% SDS at 60°C, twice for 15 min each with 0.3M NaCl, 0.5% SDS at 60°C and once for 15 min with 0.03M NaCl, 0.5% SDS at 55°C. Membranes were again placed in heat-sealed bags and exposed to Kodak RP X-ray film to 16-48 h at -70°C. The X-ray film was developed following the manufacturer's protocols. Colonies giving strong or weak signals were picked and cultured in 2xYT media supplemented with 70 µg/ml ampicillin. Plasmid DNA was isolated from these cultures using the alkaline lysis method (Sambrook et al., pp. 1.25-1.28) and analyzed by restriction digest, Southern hybridization (Sambrook et al., pp. 9.38-9.44) and PCR analysis (Sambrook et al., pp. 14.18-14.19).

Clones carrying the *bgl1* gene were identified by colony lift hybridization of the pUC119-*Hind* III library (Example 2) with a 1.0 kb *bgl1* probe prepared using oligonucleotide primers designed to amplify bp 462-1403 of the published *bgl1* sequence (Barnett et al.). A *bgl1* clone, pJEN200, was isolated containing a 6.0 kb *Hind* III fragment corresponding to the promoter, structural gene and termination sequences. Clones carrying the *cbh1* gene were identified by colony lift hybridization of the pUC119-*Bam*HI library with a 0.7 kb *cbh1* probe prepared using oligonucleotide primers designed to amplify bp 597-1361 of the published *cbh1* sequence (Shoemaker, Schweikart, Ladner, Gelfand, Kwok, Myambo and Innis, "Molecular cloning of exo-cellobiohydrolase 1 derived from *Trichoderma*

reesei strain L27", Bio/Technology 1: 691-696, 1983 hereafter referred to as Shoemaker et al.). A *cbh1* clone, pCOR132, was isolated containing a 5.7 kb *Bam*HI fragment corresponding to the promoter (4.7 kb) and 1 kb of the *cbh1* structural gene. From this, a 2.5 kb *Eco*RI fragment containing the *cbh1* promoter (2.1 kb) and 5' end of the *cbh1* coding region (0.4 kb) was subcloned into pUC119 to generate pCB152. Clones carrying the *cbh2* gene were identified by colony lift hybridization of the pUC119-*Eco*RI library with a 1.5 kb *cbh2* probe prepared using oligonucleotide primers designed to amplify bp 580-2114 of the published *cbh2* sequence (Chen, Gritzali and Stafford, "Nucleotide sequence and deduced primary structure of cellobiohydrolase II from *Trichoderma reesei*", Bio/Technology 5: 274-278, 1987, hereafter referred to as Chen et al.). A *cbh2* clone, pZUK600, was isolated containing a 4.8 kb *Eco*RI fragment corresponding to the promoter (600 bp), structural gene (2.3 kb) and terminator (1.9 kbp).

EXAMPLE 4

Cloning of *T. reesei* M2C38 *cbh1* terminator, xylanase II (*xln2*) gene, phosphoglycerate kinase promoter (*pgk p*).

Digoxigen-11-dUTP labelled probes were prepared from PCR amplified coding regions of the *cbh1*, *xln2* and *pgk* genes by random prime labeling using the DIG Labeling and Detection kit (Boehringer Mannheim) and following the manufacturer's protocols. Genomic clones containing the *cbh1*, *xln2* and *pgk* genes were identified by plaque-lift hybridization of the lambdaDASH library. For each gene of interest, 1×10^4 clones were transferred to Nytran® (Schleicher and Schull) nylon membranes. The phage particles were lysed and the phage DNA denatured by placing the membranes plaque-side up on blotting paper (VWR238) saturated with 0.5 M NaOH, 1 M NaCl for 5 min; the membranes were then neutralized by placing them plaque-side up onto blotting paper (VWR238) saturated with 1.5 M Tris, pH 7.5 plus 1 M NaCl for 5 min; the membranes were allowed to air-dry for 30 min and the DNA was then fixed to

the membranes by baking at 80°C for 2 h. The membranes were prehybridized in heat-sealed bags in a solution of 6X SSPE, 5X Denhardt's, 1% SDS plus 100 µg/ml denatured, sheared salmon sperm DNA at 65°C for 2 h. The membranes were then hybridized in heat-sealed bags in the same solution containing 50 µg/ml denatured, sheared salmon sperm DNA and 0.5 µg of digoxigenin-dUTP labelled probes at 65°C overnight. The membranes were washed twice for 15 min in 2X SSPE, 0.1% SDS at RT, twice for 15 min in 0.2X SSPE, 0.1% SDS at 65°C and once for 5 min in 2X SSPE. Positively hybridizing clones were identified by reaction with an anti-digoxigenin/alkaline phosphatase antibody conjugate, 5-bromo-4-chloro-3-indoyl phosphate and 4-nitro blue tetrazolium chloride (Boehringer Mannheim) following the manufacturer's protocol. Positively hybridizing clones were purified further by a second round of screening with the digoxigenin-dUTP labeled probes. Individual clones were isolated and the phage DNA purified as described in Sambrook, et al. (1989) pp. 2.118-2.121 with the exception that the CsCl gradient step was replaced by extraction with 1 volume of phenol:chloroform:isoamyl alcohol (25:24:1) and 1 volume of chloroform:isoamyl alcohol (24:1). The DNA was precipitated with 0.1 volume of 3M sodium acetate, and pH 5.2 and 2.5 volumes cold 95% ethanol. The precipitated phage DNA was washed with 0.5 ml cold 70% ethanol, air-dried and resuspended in 50 µl 10 mM Tris, 1 mM EDTA pH8.0. Restriction fragments containing the genes of interest were identified by restriction digests of the purified phage DNA and Southern blot hybridization (Sambrook et al., pp. 9.38-9.44) using the same digoxigenin-dUTP labeled probes used to screen the lambdaDASH library. The membranes were hybridized and positively hybridizing fragments visualized by the same methods used for the plaque lifts. Once the desired restriction fragments from each lambda DASH clone were identified, the restriction digests were repeated, the fragments were resolved on a 0.8% agarose gel in TAE and the desired bands excised. The DNA was eluted from the gel slices using the Sephaglas BandPrep Kit (Pharmacia) following the manufacturer's protocol.

Clones carrying the *cbh1* gene were identified by colony lift hybridization of the lambdaDASH library (example 2) with a *cbh1* probe comprising bp 45-2220 of the published *cbh1* sequence (Shoemaker et al.). A 1.8 kb *Bam*HI fragment containing the 3' end of the *cbh1* coding region (0.5 kb) and the *cbh1* terminator (1.3 kb) was isolated by restriction digestion of phage DNA purified from a lambdaDASH *cbh1* clone. This fragment was subcloned into the *Bam*HI site of the *E. coli* plasmid vector pUC119 to generate the plasmid pCB1Ta. Clones carrying the *xln2* gene were identified by colony lift hybridization of the lambdaDASH library (example 2) with a *xln2* probe comprising bp 100-783 of the published *xln2* sequence (Saarelainen, Paloheimo, Fagerstrom, Suominen and Nevalainen, "Cloning, sequencing and enhanced expression of the *Trichoderma reesei* endoxylanase II (pI 9) gene *xln2*", Mol. Gen. Genet. 241: 497-503, 1993, hereafter referred to as Saarelainen et al.). A 5.7 kb *Kpn*I fragment containing the promoter (2.3 kb), coding region (0.8 kb) and terminator (2.6 kb) the *xln2* gene was isolated by restriction digestion of phage DNA purified from a lambdaDASH *xln2* clone. This fragment was subcloned into the *Kpn*I site of pUC119 to generate the plasmid pXYN2K-2. Clones carrying the *pgk* gene were identified by colony lift hybridization of the lambdaDASH library (Example 2) with a *pgk1* probe comprising bp 4-1586 of the published *pgk* sequence (Vanhanen, Penttila, Lehtovaara and Knowles, "Isolation and characterization of the 3-phosphoglycerate kinase gene (*pgk*) from the filamentous fungus *Trichoderma reesei*", Curr. Genet. 15: 181-186, 1989). A 5.0 kb *Eco*RI fragment containing the promoter (2.9 kb), coding region (1.6 kb) and terminator (0.5 kb) of the *pgk* gene was isolated by restriction digestion of phage DNA purified from a lambdaDASH *pgk* clone. This fragment was subcloned into the *Eco*RI site of pUC119 to generate the plasmid pGK5.0.

EXAMPLE 5

Construction of β -glucosidase overexpression vector pCBG1-TV

This Example describes the construction of a vector containing the *Trichoderma* cellobiohydrolase I promoter and secretion signal and the mature beta-glucosidase coding region.

A DNA fragment containing the *bgl1* coding region minus the β -glucosidase secretion signal (bp 474-2679) was amplified by PCR from the pJEN200 template using primers homologous to the published *bgl1* sequence containing either an *SphI* site 5' to Val32 of the encoded β -glucosidase or a *KpnI* site 3' to the *bgl1* stop codon using *Pwo* polymerase. This amplified fragment was digested with *SphI* and *KpnI* and inserted into pCB219N digested with *SphI* and *KpnI* to generate pBgstrf. To make pCB219N, a *cbh2* terminator fragment was amplified from the pZUK600 template using a primer homologous to bp 2226-2242 of the published 3' untranslated region of the *cbh2* gene (Chen et al., 1987) containing a *KpnI* site at the 5' end and the pUC forward primer (Cat. No. 1224, New England Biolabs) which anneals downstream of the *EcoRI* site at the 3' end of *cbh2* in pZUK600. This fragment was digested at the engineered *KpnI* and *EcoRI* sites and inserted into the corresponding sites of pUC119 to generate pCB219. An *EcoRI*-*NotI* adaptor (Cat. No. 35310-010, Gibco/BRL) was inserted into the unique *EcoRI* site of pCB219 to generate pCB219N. A fragment containing the *cbh1* promoter and secretion signal was amplified from pCB152 using a *cbh1* specific primer (bp 249-284 of the published *cbh1* sequence, Shoemaker et al., 1983) containing a *SphI* site 3' to Ser19 of the encoded CBH1 and pUC forward primer (Cat. No. 1224, New England Biolabs) which anneals upstream of the *EcoRI* site at the 5' end of *cbh1* in pCB152. The *cbh1* promoter + secretion signal PCR product was digested with *SphI* and *EcoRI* and inserted into the corresponding sites in pBR322L (a derivative of pBR322 in which the region between the *SphI* and *SalI* sites was replaced with an *SphI*-*NotI*-*SalI* linker) to generate pBR322LCS. To make the expression cassette, the *bgl1* coding region and *cbh2* terminator were isolated

from pBgstrf as a 4.1 kb *SphI*/*NotI* fragment and inserted into pBR322LCS digested with *SphI* and *NotI*. In order to maintain the correct reading frame at the juncture of the *cbh1* secretion signal and the mature β -glucosidase, the resultant plasmid, pCBGstrf, was linearized at the unique *SphI* site and the *SphI* site was blunted with T4 DNA polymerase. The resulting plasmid, pCBG1, was then further modified by conversion of the unique *NotI* site at the 3' end of the *cbh2* terminator to a unique *XhoI* site by the addition of *XhoI* linkers (Cat. No. 1073, New England Biolabs). The final plasmid, pCBG1-*Xho*, is the expression cassette plasmid.

The *E. coli* hygromycin phosphotransferase gene (*hph*) used as a selectable marker for *T. reesei* was amplified with *Pwo* polymerase from the plasmid pVU1005 (Van den Elzen, Townsend, Lee and Bedbrook, "A chimaeric hygromycin resistance gene as a selectable marker in plant cells", Plant Mol. Biol. 5: 299-302, 1989). The primers were designed to introduce *SphI* and *KpnI* sites at the 5' and 3' ends of the *hph* coding region (bp 211-1236 of the published *hph* sequence, Gritz and Davies, "Plasmid-encoded hygromycin b resistance: the sequence of hygromycin B phosphotransferase gene and its expression in *Escherichia coli* and *Saccharomyces cerevisiae*" Gene 25: 179-188, 1983), respectively. The PCR product was digested with *SphI* and *KpnI* and inserted into the corresponding sites in the polylinker region of pUC119. The resulting plasmid, pHPT100 was used as the starting plasmid for the construction of the selection cassette. Two new linker regions were introduced into this plasmid to facilitate the cloning of the promoter and terminator fragments. A *HindIII*-*XbaI*-*XhoI*-*SphI* linker was inserted between the *HindIII* and *SphI* sites as well as a *KpnI*-*NotI*-*SacI* linker which was inserted between the *KpnI* and *SacI* sites of pUC119 polylinker remaining in pHPT100. This construct was designated as pHPT102. The primers used to amplify the *pgk* promoter (Vanhanen, Saloheimo, Ilmen, Knowles and Penttila, "Promoter structure and expression of the 3-phosphoglycerate kinase gene (*pgk1*) of *Trichoderma reesei*", Gene 106: 129-133, 1991) were designed to introduce an *XhoI* site and a *SphI* site at positions -970

and +1 of the promoter respectively. These sites were subsequently used to insert the *pgk* promoter into the *Xho*I and *Sph*I sites of pHPT102 to generate pHPT115. A 1.3 kb *cbh*1 terminator fragment was amplified with *Pwo* polymerase from pCB1Ta using a primer annealing to the 3' untranslated region of *cbh*1 (bp 1864-1899 of the published *cbh*1 sequence) containing a *Kpn*I site at bp 1877-1882 and the pUC reverse primer (Cat. No., 18432-013, Gibco/BRL) which anneals downstream of the *Eco*RI site at the 3' end of the *cbh*1 terminator in pCB1Ta. The *cbh*1 terminator PCR product was digested with *Kpn*I and inserted into the unique *Kpn*I site of pHPT115 to generate the selection cassette plasmid pHPT136.

To make the transformation vector, the expression cassette from pCBG1-Xho was isolated as a 5.6 kb *Xba*I/*Xho*I fragment and inserted between the unique *Xba*I and *Xho*I sites upstream of the selection cassette of pHPT136. The final transformation vector, pCBG1-TV, as depicted in Figure 1, was introduced as a circular plasmid into *T. reesei* M2C38 via microprojectile bombardment as described below in Example 9.

EXAMPLE 6

Construction of β -glucosidase overexpression vector pXBG1-TV

This Example describes the construction of a vector containing the *Trichoderma* xylanase II promoter and secretion signal, and the mature beta-glucosidase coding region.

The β -glucosidase coding region (bp 474-2680) was amplified with *Pwo* polymerase from the genomic *bgl*1 clone pJEN200 using primer to insert a *Xba*I site directly upstream of bp 474 in the published *bgl*1 sequence (Barnett, et al.) and a *Kpn*I site directly downstream of bp 2680. The blunt ended PCR product was inserted into the *Sma*I site of pUC118 to generate the plasmid designated as pBGm.s. Since the *Xba*I site was engineered to be immediately upstream of the

start of the mature β -glucosidase at Val32, the cloned fragment did not include the β -glucosidase secretion signal. The plasmid pBGm.s was digested with *Xba*I and *Kpn*I and the 2.2 kb fragment containing the *bgl*I coding region minus the secretion signal was isolated and inserted into the *Xba*I and *Kpn*I sites upstream of the *cbh*2 terminator in the plasmid pCB219N (described in Example 5, above) to yield the plasmid pBG2X. A 2.3 kb fragment containing the promoter and secretion signal of the *xln*2 gene (bp -2150 to +99 where +1 indicates the ATG start codon) was amplified with *Pwo* polymerase from the genomic *xln*2 subclone pXYN2K-2 using a *xln*2-specific primer containing a *Nhe*I site directly downstream of bp103 of the published *xln*2 sequence (Saarelainen et al.) and the pUC reverse primer (Cat. No. 18432-013, Gibco/BRL) which anneals upstream of the *Kpn*I site at the 5' end of the *xln*2 gene. This *xln*2 PCR product was digested with *Eco*RI (which was amplified as part of the pUC119 polylinker from pXYN2K-2) and *Nhe*I and inserted into the plasmid pBR322L (described in example 5 above) to generate pBR322LXN. The *Eco*RI site of pBR322LXN was then blunted with Klenow, and *Spe*I linkers (Cat. No. 1086, New England Biolabs) were added to generate pBR322SpXN. The pBG2X plasmid was cut with *Xba*I and *Not*I and a 4.2 kb fragment, containing the *bgl*I coding region followed by the *cbh*2 terminator, was isolated. This fragment was inserted into the plasmid pBR322SpXN cut with *Nhe*I and *Not*I (*Nhe*I and *Xba*I have compatible overhangs). This cloning resulted in the fusion of the xylanase secretion signal directly to the mature β -glucosidase creating the complete expression cassette pXBG-2.

The *cbh*1 terminator in the selection cassette plasmid pHPT136 described in Example 5, above, was replaced with a 2.6 kb *Kpn*I fragment containing the *xln*2 transcriptional terminator. The *xln*2 terminator was amplified with *Pwo* polymerase from the genomic subclone pXYN2K-2 using a primer to introduce a *Kpn*I site directly downstream of bp 780 of the published *xln*2 sequence (Saarelainen et al.) and the pUC forward primer (Cat. No. 18431-015, Gibco/BRL) which anneals downstream of the 3' end of the *xln*2 gene in pXYN2K-2. The

xln2 terminator PCR product was digested with *KpnI* and ligated to a 5.1 kb *KpnI* fragment from pHPT136 containing the *pgk* promoted-*hph* gene in pUC119 to generate the selection cassette plasmid pHPT136X.

The construction of the transformation vector involved the insertion of the expression cassette directly upstream of the *pgk* promoter from the selection cassette. The expression cassette plasmid pXBG2 was digested with *NotI*, the ends were made blunt using Klenow, and then digested with *SpeI*. The selection cassette pHPT136X was prepared in a similar manner by digestion with *XhoI*, followed by the fill in reaction to create the blunt ends and then a digestion with *XbaI*. A blunt-sticky ligation of these two fragments was performed to ligate the 6.5 kb *SpeI*/blunted *NotI* fragment from pXBG2 into the *XbaI*/blunted *XhoI* fragment of pHPT136X (*SpeI* and *XbaI* have compatible overhangs). The final transformation vector, pXBG-TV, as depicted in Figure 2, was linearized at its unique *NotI* prior to transformation of *T. reesei* M2C38 via microprojectile bombardment, as described below in Example 9.

EXAMPLE 7

Construction of β -glucosidase overexpression vector pC/XBG(*XbaI*)-TV

This Example describes the construction of a vector containing the *Trichoderma* cellobiohydrolase 1 promoter, the xylanase II secretion signal and the mature beta-glucosidase coding region.

This Example was carried out to test the combined effects of the *cbh1* promoter and *xln2* secretion signal on *bgl* expression. A 1.2 kb *HindIII* fragment comprising bp -1399 to -204 of the *cbh1* promoter was amplified by PCR using the *cbh1* promoter-containing plasmid pBR322LCS (Example 5) as a template in order to insert a unique *XbaI* site at bp -1393 to -1388. This modified *cbh1* promoter fragment was digested with *HindIII* and was used to replace bp -1400 to

-121 of the *xln2* promoter in pXBG1 (Example 6) to generate the new expression cassette plasmid pC/XBG1. The 6.4 kb expression cassette from pC/XBG1 was isolated by digestion with *NotI* followed by blunting of the *NotI* site with Klenow fragment and subsequent digestion with *SpeI*. This fragment was then inserted by blunt/sticky ligation upstream of the *hph* selection cassette in pHPT136X which had been digested with *XhoI*, blunted at the *XhoI* site with Klenow and digested with *XbaI*. The final transformation vector, pC/XBG(*XbaI*)-TV (Accession No. 209613, deposit date February 3, 1998, American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852 USA), as shown in Figure 3, was linearized at the unique *XbaI* and *NotI* sites at the 5' end of the *cbh1* promoter and the 3' end of the *xln2* terminator prior to transformation of *T. reesei* M2C38 via microprojectile bombardment, as described below in Example 9.

EXAMPLE 8

Southern blots of genomic DNA isolated from *T. reesei* strains RutC30 and M2C38

Genomic DNA was isolated from each strain as described in Example 1. For Southern blots, 1 µg of DNA was digested with 3-10 units of restriction endonuclease at 37°C for at least 2 hours and the digestion products resolved on a 0.8% agarose gel in 0.04 M Tris-acetate, 1 mM EDTA. DNA was transferred by nylon membranes (Boehringer Mannheim) by capillary transfer (Sambrook et al., pp. 9.38-9.44). In Figures 4 and 5, lanes 2, 4, 6, 8, 10 and 12 contain digested M2C38 DNA and lanes 3, 5, 7, 9, 11 and 13 contain digested RutC30 DNA. The restriction endonucleases used were *BamHI* (lanes 2 and 3), *EcoRI* (lanes 4 and 5), *XbaI* (lanes 6 and 7), *Hind III* (lanes 8 and 9), *SstI* (lanes 10 and 11) and *KpnI* (lanes 12 and 13). In both figures, lane 1 contains lambda *HindIII* molecular size standards (Gibco/BRL, cat. no. 15612-013) and lane 14 contains 1 ng of unlabeled fragment used to make the probe. Southern blots were hybridized with a digoxigenin-11-dUTP labelled random-primed probe prepared using the DIG

Labeling and Detection Kit (Boehringer Mannheim). The template for the probe used in Figure 4 was a 2.3 kb fragment comprising the *T. reesei xln2* promoter and secretion signal (Saarelainen et al.). The template for the probe used in Figure 5 was a 2.1 kb fragment comprising bp 574-2679 of the *T. reesei bgl1* mature coding region (Barnett, et al.). After post-hybridization washes, dig-dUTP complexes were visualized by incubation with an anti-digoxigenin alkaline phosphatase conjugate (Boehringer Mannheim) followed by reaction with 5-bromo-4-chloro-3-indoyl phosphate and 4-nitro blue tetrazolium chloride (Boehringer Mannheim).

EXAMPLE 9

Transformation of *T. reesei* RutC30, M2C38 and BTR48 via microprojectile bombardment

The Biolistic PDS-1000/He system (BioRad; E.I. DuPont de Nemours and Company) was used to transform spores of *T. reesei* strains RutC30, M2C38 and BTR48, and all procedures were performed as recommended by the manufacturer. M-10 tungsten particles (median diameter of 0.7 μ m) were used as microcarriers. The following parameters were used in the optimization of the transformation: a rupture pressure of 1100 psi, a helium pressure of 29 mm Hg, a gap distance of 0.95 cm, a macrocarrier travel distance of 16 mm, and a target distance of 9 cm. Plates were prepared with 1×10^6 spores on Potato Dextrose Agar media (PDA). Bombarded plates were incubated at 28°C. Four hours post-bombardment, spores are subjected to primary selection by the overlaying of selective PDA media supplemented with 80 units/ml of HygB. The bombardment plates are incubated at 28°C. Transformants can be observed after 3-6 days growth; however, further incubation is necessary to achieve sporulation.

After sporulation has occurred, a secondary selection process is performed to isolate individual transformants. Spores are collected from the plate with an

inoculating loop and resuspended in sterile water. This suspension is then filtered through a sterile syringe plugged with glass microfibers. This allows the passage of spores while retaining unwanted mycelia. A determination of the concentration of spores in this suspension is required and subsequent dilutions are plated onto PDA plates supplemented with 0.75% Oxgall (Difco) and HygB (40 units/mL) to obtain 20-50 spores per plate. The Oxgall acts as a colony restrictor, thereby allowing the isolation of individual colonies on these secondary selection plates. Isolated colonies can be observed after 2-3 days.

EXAMPLE 10

Southern blot analysis of genomic DNA isolated from *T. reesei* strains RutC30, RC300, RC-302, M2C38, RM4-300, R4-301, RM4-302, BTR48, and RB4-301

Genomic DNA was isolated from each strain as described in Example 1. For Southern blots, 1 µg of DNA was digested with 3-10 units of *Kpn*I or *Xba*I at 37°C for at least 2 hours and the digestion products resolved on a 0.8% agarose gel in 0.04 M Tris-acetate, 1 mM EDTA. DNA was transferred by nylon membranes (Boehringer Mannheim) by capillary transfer (Sambrook et al., pp. 9.38-9.44). Southern blots were hybridized with a digoxigenin-11-dUTP labelled random-primed probe prepared using the DIG Labeling and Detection Kit (Boehringer Mannheim). The template was a 1.3 kb *Eco*R1-*Bgl* II fragment comprising bp1215-2464 of the published *bgl*I sequence (Barnett et al.). After post-hybridization washes, dig-dUTP complexes were visualized by incubation with an anti-digoxigenin alkaline phosphatase conjugate (Boehringer Mannheim) followed by reaction with the chemiluminescent reagent CSPD (Boehringer Mannheim) and exposure to X-ray film (Kodak). The results are summarized in Table 1.

TABLE 1
bgl1 copy number in parental and recombinant *T. reesei* strains

Strain	Host	Promoter	Secretion signal	Vector	native <i>bgl1</i> gene	# <i>bgl1</i> vectors	total # <i>bgl1</i> genes
RutC30	Same	<i>bgl1</i>	<i>bgl1</i>	none	Present	0	1
RC-300	RutC30	<i>cbh1</i>	<i>cbh1</i>	pCBG1-TV	Present	1	2
RC-302	RutC30	<i>cbh1</i>	<i>xln2</i>	pC/XBG1-TV	Absent	1	1
M2C38	Same	<i>bgl1</i>	<i>bgl1</i>	None	Present	0	1
RM4-300	M2C38	<i>cbh1</i>	<i>cbh1</i>	pCBG1-TV	Absent	2	2
RM4-301	M2C38	<i>xln2</i>	<i>xln2</i>	pXBG1-TV	Present	2	3
RM4-302	M2C38	<i>cbh1</i>	<i>xln2</i>	pC/XBG1-TV	Present	2	3
BTR48	Same	<i>bgl1</i>	<i>bgl1</i>	None	Present	0	1
RB4-301	BTR48	<i>xln2</i>	<i>xln2</i>	pXBG1-TV	Absent	2	2

EXAMPLE 11

Production of β -glucosidase in liquid cultures

This Example describes the methods used to determine the amount of beta-glucosidase enzyme produced by a *Trichoderma* strain.

Individual colonies of *Trichoderma* are transferred to PDA plates for the propagation of each culture. Sporulation is necessary for the uniform inoculation of shake flasks which are used in testing the ability of the culture to produce the beta-glucosidase and cellulase. The culture media is composed of the following:

Component	g/L
(NH ₄) ₂ SO ₄	6.35
KH ₂ PO ₄	4.00
MgSO ₄ ·7H ₂ O	2.02
CaCl ₂ ·2H ₂ O	0.53
CSL	6.25
CaCO ₃	10.00
Carbon sources**	5-10
Trace elements*	1 mL/L

*Trace elements solution contains 5 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$; 1.6 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$; 1.4 g/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$.

** 5 g/l glucose plus 10 g/l Solka floc (when the *cbh1* or other cellulase promoter is used), 10 g/l xylan (when the *xln2* promoter is used) or other carbon source compatible with the promoter directing the expression of the beta-glucosidase. The carbon source can be sterilized separately as an aqueous solution at pH 2 to 7 and added to the remaining media.

The liquid volume per 1-liter flask is 150 mL, the initial pH is 5.5 and each flask is sterilized by steam autoclave for 30 minutes at 121°C prior to inoculation.

For both untransformed (i.e., native) and transformed cells, spores are isolated from the PDA plates as described in Example 9 and $1-2 \times 10^6$ spores are used to inoculate each flask. The flasks are shaken at 200 rpm at a temperature of 28°C for a period of 6 days. The filtrate containing the secreted protein was collected by filtration through GF/A glass microfibre filters (Whatman). The protein concentration is determined using the Bio-Rad Protein Assay (Cat. No. 500-0001) using *Trichoderma* cellulase as a standard. Beta-glucosidase activity is determined as described in Example 16.

Transformants were screened for the ability to produce at least 10-fold more beta-glucosidase (in IU/mg) than the untransformed host strain as determined by the IU/ml of beta-glucosidase activity of the culture filtrate divided by the protein concentration (in mg/ml) of the culture filtrate.

EXAMPLE 12

Production of beta-glucosidase by *T. reesei* strains RutC30, RC-300, and RC-302 using Solka floc carbon source

Based on previous successes using the *cbh1* promoter and secretion signal to overexpress proteins in *Trichoderma*, the mature beta-glucosidase coding region was placed downstream of the *cbh1* promoter and secretion signal in the genetic construct shown in Figure 1 and described in Example 5 (pCBG1-TV). The vector was introduced into *T. reesei* RutC30 by particle bombardment (Example 9) and the resulting transformant RC-300, produced 7 times more beta-glucosidase activity than the parental strain (Table 2). This 7-fold increase resulted from the incorporation of one copy of the transformation vector into the host chromosomes (Example 10, Table 1). The larger increase in beta-glucosidase activity obtained from one copy of a construct in which beta-glucosidase is expressed using the *cbh1* promoter and secretion signal suggests that this strategy is better than that employed by Barnett et al. and Fowler et al. which resulted in only a 5-fold increase in beta-glucosidase activity from 10-15 copies of a construct in which beta-glucosidase is expressed from its own promoter and secretion signal. However, the resulting 7-fold increase in beta-glucosidase activity was still not sufficient to alleviate the shortage of beta-glucosidase for cellulose hydrolysis.

The untransformed *T. reesei* strain RutC30 was transformed by particle bombardment (Example 9) with a genetic construct from the vector pC/XBG(Xba1)-TV encoding the mature *T. reesei* beta-glucosidase enzyme linked to the *T. reesei* xylanase II secretion signal.

The untransformed strain RutC30 and the resulting transformed strain from this host, RC-302, were cultured using the procedures of Example 11 with 10 g/L Solka floc and 5 g/L glucose as carbon sources. The results are shown in Table 2.

The untransformed strain produced 0.14 IU of beta-glucosidase per mg protein.

The transformant RC-302 with the CBH1 promoter and xylanase II secretion signal produced 19 IU/mg of beta-glucosidase. This represents about a 136-fold improvement over the untransformed strain, which is very significant for a cellulose-to-ethanol process.

The transformant RC-302 with the CBH1 promoter and xylanase II secretion signal produced about 19 times more beta-glucosidase activity than the best RutC30 transformant with the CBH1 promoter and CBH1 secretion signal.

TABLE 2

Production of β -glucosidase in *T.reesei* strains RutC30, RC-300, and RC-302 in 150 ml flask cultures

Strain	promoter	Secretion signal	β -g (IU/mg)
RutC30	bgl1	bgl1	0.14
RC-300	cbh1	cbh1	1.0
RC-302	cbh1	xln2	19

EXAMPLE 13

Production of beta-glucosidase by strains M2C38 and RM4-302 using Solka floc carbon source

The vector pCBG1-TV, in which the beta-glucosidase is expressed from the CBH1 promoter and secretion signal (Figure 1 and Example 5), was introduced into *T. reesei* M2C38 by particle bombardment (Example 9). The resulting transformant RM4-300 produced about 7-12 times more beta-glucosidase activity than the parental strain (Table 3).

The untransformed *T. reesei* strain M2C38 was transformed by particle bombardment (Example 9) with a genetic construct from the vector pC/XBG(Xba1)-TV encoding the mature *T. reesei* beta-glucosidase enzyme linked to *T. reesei* xylanase II secretion signal.

The untransformed strain M2C38 and the transformed strain from this host, RM4-302, were cultured using the procedures of Example 11 with 10 g/L Solka floc and 5 g/L glucose as carbon sources. The results are shown in Table 3.

The untransformed strain produced 0.35 IU of beta-glucosidase per mg protein.

The transformant RM4-302 with the CBH1 promoter and xylanase II secretion signal produced 14.1 IU/mg of beta-glucosidase. This represents about a 40-fold improvement over the untransformed strain, which is very significant for a cellulose-to-ethanol process.

The transformant RM4-302 with the CBH1 promoter and xylanase II secretion signal produced about 3 times more beta-glucosidase activity than the transformant with the CBH1 promoter and CBH1 secretion signal. This is a significant difference, as the CBH1 promoter and secretion signal did not lead to sufficient production of beta-glucosidase to completely suppress cellobiose production in hydrolysis.

TABLE 3

Production of β -glucosidase in *T. reesei* strains M2C38, RM4-300, and RM4-302 in 150 ml flask cultures

Strain	promoter	Secretion signal	β -g (IU/mg)
M2C38	bgl1	bgl1	0.35
RM4-300	cbh1	cbh1	4.5
RM4-302	cbh1	xln2	14.1

EXAMPLE 14

Production of beta-glucosidase by *T. reesei* strains M2C38 and RM4-301
using xylan carbon source

The untransformed *T. reesei* strain M2C38 was transformed by particle bombardment (Example 9) with a genetic construct from the vector pXBG1-TV encoding the mature *T. reesei* beta-glucosidase linked to the xylanase promoter and secretion signal.

The untransformed strain M2C38 and a transformed strain from this host, RM4-301, were cultured using the procedures of Example 11 with 5 g/L glucose and 10 g/L xylan as the carbon source. The results are shown in Table 4.

The untransformed strain produced 0.16 IU of beta-glucosidase per mg protein. The transformant RM4-301 with the xylanase II promoter and xylanase II secretion signal produced 20.4 IU/mg of beta-glucosidase. This represents about a 127-fold improvement over the untransformed strain, which is very significant for a cellulose-to-ethanol process.

TABLE 4

Production of β -glucosidase in *T. reesei* strains M2C38 and RM4-301
with xylan in 150 ml flask cultures

Strain	promoter	Secretion signal	β -g (IU/mg)
M2C38	bgl1	bgl1	0.16
RM4-301	xln2	xln2	20.4

EXAMPLE 15

Production of beta-glucosidase by strains BTR-48 and RB48-301
using Solka floc carbon source

The untransformed *T. reesei* strain BTR48 was transformed by particle bombardment with a genetic construct from the vector pXBG1-TV encoding the mature *T. reesei* beta-glucosidase linked to the xylanase promoter and secretion signal.

The untransformed strain BTR-48 and a transformed strain from this host, RB48-301, were cultured using the procedures of Example 11 with 5 g/L glucose and 10 g/L Solka floc as the carbon sources. The results are shown in Table 5.

The untransformed strain produced 0.16 IU of beta-glucosidase per mg protein. The transformant RB48-301 with the xylanase II promoter and xylanase II secretion signal produced about 21.9 IU/mg of beta-glucosidase. This represents about a 136-fold improvement over the untransformed strain, which is very significant for a cellulose-to-ethanol process.

TABLE 5

Production of β -glucosidase in *T.reesei* strains BTR48 and RB48-301 with Solka floc in 150 ml flask cultures

Strain	promoter	Secretion signal	β -g (IU/mg)
BTR48	bgl1	bgl1	0.16
RB48-301	xln2	xln2	21.9

EXAMPLE 16

Measurement of beta-glucosidase activity of an enzyme mixture

The beta-glucosidase activity of an enzyme is measured using the procedures of Ghose, "Measurement of Cellulase Activities," Pure and Appl. Chem., 59:257-268 (1987), as follows. The sample of enzyme is diluted to several concentrations in 50 mM sodium citrate buffer, pH 4.8, to a volume of 0.5 ml. A convenient range

of dilutions is 3 to 24 times the estimated activity of the sample. For example, a 10 unit/ml sample should be diluted 1:30 to 1:240. Regardless of the dilutions used, a sample of 0.5 ml of the citrate buffer is added to each enzyme tube. The substrate is prepared as 15 mM (5.13 g/L) cellobiose. The dilute enzyme stocks and the substrate are separately preheated to 50°C for 5 minutes, then a 0.5 ml aliquot of the substrate is added to each tube with enzyme. The test tubes are incubated for 30 minutes at 50°C. The reaction is terminated by immersing each tube in a boiling water bath for 5 minutes. The tubes are then vortex mixed, and the amount of sugar produced by each sample of enzyme is measured on a YSI glucose analyzer, taking into account the small background from the enzyme.

A unit of beta-glucosidase activity is defined as the number of micromoles of glucose produced per minute. The activity is calculated based on Equation 1 using the average value from each of the dilutions which produces 0.15 to 1.5 mg/ml of glucose.

$$A = C \cdot G \cdot D \quad (1)$$

where A = activity, beta-glucosidase units/ml
(or micromoles glucose/ml/min)
C = 16.7 micromoles/mg/min
G = glucose produced, mg/ml
D = enzyme dilution, dimensionless

EXAMPLE 17

Cellulose hydrolysis

The purpose of this experiment was to demonstrate the effectiveness of the beta-glucosidase made by the transformed *Trichoderma* in enhancing the hydrolysis of cellulose.

The enzymes used for this study were Iogen Cellulase, a commercial cellulase enzyme of Iogen Corporation, and the product of RM4-302 grown in a 30-liter

fermentation vessel using the procedures described in Example 11, with twice the media concentration levels listed in that Example. The enzyme concentration was increased by ultrafiltration across an Amicon 10,000 MWCO membrane and normalized to the same cellulase activity as Iogen Cellulase. The activities of these two enzymes are shown in Table 6.

TABLE 6
Enzyme activities used in cellulose hydrolysis study

Enzyme	Beta-glucosidase IU/ml	Cellulase FPU/ml	BG IU/mg @ 10 FPU/g
Iogen Cellulase	112	140	8.0
RM4-301	1170	140	83.6

The cellulose used for this study was pretreated oat hulls, prepared as per the procedures of Foody, et al, Improved Pretreatment Process for Conversion of Cellulose to Fuel Ethanol, US patent application filed June 9, 1997, Example 6.

Samples of pretreated oat hull cellulose of 0.5 grams were added to 25 ml flasks with 49.5 grams of 0.05 molar sodium citrate buffer, pH 4.8 .

The enzymes were added to the flask in an amount corresponding to 10 FPU per gram of cellulose. The resulting beta-glucosidase dosages are listed in Table 6.

In both cases, the flasks were shaken at 250 RPM and maintained at 50°C for 24 hours. At this time, samples were taken, filtered to remove insoluble cellulose, and analyzed for glucose and cellobiose concentration using standard Dionex pulsed-amperometric HPLC carbohydrate analysis methods. The results are listed in Table 7.

Iogen Cellulase, the conventional *Trichoderma* cellulase, converted only 45% of the cellulose to glucose. This is unacceptably low for an ethanol process. The accumulation of cellobiose was significant, representing 13% of the cellulose.

The cellulase with enhanced beta-glucosidase performed much better. The cellulose conversion to glucose reached 84%. The reason for this excellent performance was that cellobiose accumulation was negligible, due to the abundance of beta-glucosidase.

TABLE 7

Hydrolysis of cellulose enhanced by high beta-glucosidase

Enzyme	Glucose (% of cellulose)	Cellobiose (% of cellulose)
Iogen Cellulase	45	13
RM4-301	84	< 1

EXAMPLE 18

Comparison of the *Trichoderma reesei xln2* and *bgl1* genes in strains RutC30 and M2C38

Southern Blot analysis was performed on M2C38 and RutC30 DNA digested with six different restriction enzymes that cut both within and outside of the regions that encode the mature beta-glucosidase and the xylanase secretion signal (Example 8) to determine if any polymorphisms exist between the two strains. As shown in Figures 4 and 5, the identical bands were found to hybridize with labelled probes prepared from M2C38 fragments encoding the mature beta-glucosidase enzyme and the xylanase II promoter plus secretion signal, indicating no polymorphisms and a high degree of DNA sequence homology in these regions between the two strains.

The probes and primers used to identify and clone the M2C38 DNA sequences necessary to make the genetic constructs described in Examples 5-7 were based on published DNA sequences of the various genes from several different *Trichoderma reesei* strains including QM9414 (*pgk*, Vanhanen et al., 1989 and *cbh2*, Chen et al.), the QM9414 derivatives VTT-D79125 (*xln2*, Saarelainen et al.) and L27 (*cbh1*, Shoemaker et al.), and the strain RL-P37 derivative strain P40 (*bgl1*, Barnett et al.). All of these strains, like M2C38, are derived from strain QM6a (Carter, Allison, Rey and Dunn-Coleman, "Chromosomal and genetic analysis of the electrophoretic karyotype of *Trichoderma reesei*: mapping of the cellulase and xylanase genes," Molecular Microbiology 6: 2167-2174, 1992).

Because RutC30 is the QM6a-derived progenitor of M2C38, the inventors are confident that the method as described in Examples 2-4, for the isolation of the gene sequences used to make the beta-glucosidase expression vectors described in Examples 5-7, will work equally well for the isolation of the same gene sequences from both M2C38 and RutC30. Based on the strain lineage described above and the Southern blot data, the inventors also have a high degree of confidence that genetic constructs prepared from RutC30 DNA will contain the identical DNA segments encoding the mature beta-glucosidase enzyme and the xylanase II secretion signal as those prepared from M2C38 DNA. Since the constructs prepared from M2C38 DNA (Examples 5-7) result in enhanced beta-glucosidase expression in both M2C38 and RutC30 (Examples 12-14), the inventors are also confident that genetic constructs made from RutC30 DNA will result in similar levels of enhancement of beta-glucosidase activity in both RutC30 and M2C38.

While the present invention has been described with respect to what is at present considered to be the preferred embodiments, it is to be understood that the invention is not limited to the disclosed embodiments. To the contrary, the invention is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims. The scope of the following claims is to be accorded the broadest interpretation so as to encompass all such modifications and equivalent formulations and functions.

WE CLAIM:

1. A genetically modified microbe for producing an increased level of beta-glucosidase, comprising in combination:
 - a microbe selected from the group consisting of *Trichoderma*, *Humicola*, *Fusarium*, *Streptomyces*, *Thermomonospora*, *Bacillus*, *Cellulomonas*, and *Aspergillus*; and
 - a genetic construct that has been introduced into said microbe, said genetic construct having a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region,wherein said genetically modified microbe produces an increased level of beta-glucosidase relative to said microbe.
2. The genetically modified microbe of Claim 1, wherein said microbe is a *Trichoderma* microbe.
3. The genetically modified microbe of Claim 2, wherein said *Trichoderma* microbe is a *Trichoderma reesei* microbe.
4. The genetically modified microbe of Claim 1, wherein said genetically modified microbe produces an increased level of beta-glucosidase of at least about 10-fold.
5. The genetically modified microbe of Claim 1, wherein said genetically modified microbe produces an increased level of beta-glucosidase of at least about 40-fold.

6. The genetically modified microbe of Claim 1, wherein said genetically modified microbe produces an increased level of beta-glucosidase of at least about 120-fold.
7. The genetically modified microbe of Claim 1, wherein said xylanase secretion signal is native to said microbe from which said genetically modified microbe is derived.
8. The genetically modified microbe of Claim 1, wherein said xylanase secretion signal comprises a xylanase secretion signal of a Family 11 xylanase gene.
9. The genetically modified microbe of Claim 8, wherein said Family 11 xylanase gene comprises a *Trichoderma* xylanase gene.
10. The genetically modified microbe of Claim 9, wherein said *Trichoderma* xylanase gene comprises a *Trichoderma* xylanase I gene or a *Trichoderma* xylanase II gene.

11. The genetically modified microbe of Claim 9, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma bglI* gene, and wherein said promoter is selected from the group consisting of *Trichoderma cbh1*, *cbh2*, *eg1*, *eg2*, *eg3*, *eg5*, *xln1*, and *xln2* promoter.
12. The genetically modified microbe of Claim 1, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
13. The genetically modified microbe of Claim 2, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
14. The genetically modified microbe of Claim 3, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
15. The genetically modified microbe of Claim 4, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.

16. The genetically modified microbe of Claim 5, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
17. The genetically modified microbe of Claim 6, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
18. The genetically modified microbe of Claim 7, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
19. The genetically modified microbe of Claim 8, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
20. The genetically modified microbe of Claim 9, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
21. The genetically modified microbe of Claim 10, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.

22. A genetic construct useful for expressing beta-glucosidase when introduced into a microbe, said genetic construct comprising:
- a promoter selected from the group consisting of *Trichoderma cbh1*, *cbh2*, *eg1*, *eg2*, *eg3*, *eg5*, *xln1*, and *xln2* promoters;
 - a xylanase secretion signal of a Family 11 xylanase gene; and
 - a mature beta-glucosidase coding region of a beta-glucosidase gene selected from the group consisting of *Trichoderma*, *Aspergillus*, *Humicola*, and *Fusarium* beta-glucosidase genes.
23. The genetic construct of Claim 22, wherein said Family 11 xylanase gene comprises a *Trichoderma* xylanase gene.
24. The genetic construct of Claim 23, wherein said *Trichoderma* xylanase gene comprises a *Trichoderma* xylanase I gene or a *Trichoderma* xylanase II gene.
25. The genetic construct of Claim 22, wherein said mature beta-glucosidase coding region comprises a mature beta-glucosidase coding region of a *Trichoderma* beta-glucosidase gene.
26. The genetic construct of Claim 25, wherein said *Trichoderma* beta-glucosidase gene comprises a *Trichoderma bglI* gene.

27. A method for producing beta-glucosidase, comprising:

transforming a microbe with a genetic construct comprising a promoter, a xylanase secretion signal, and a mature beta-glucosidase coding region, to create a genetically modified microbe; and

using the genetically modified microbe to produce an increased level of beta-glucosidase relative to the microbe prior to being transformed.

28. The method of Claim 27, wherein said transforming step comprises transforming a microbe selected from the group consisting of *Trichoderma*, *Humicola*, *Fusarium*, *Streptomyces*, *Thermomonospora*, *Bacillus*, *Cellulomonas*, and *Aspergillus*, to create a genetically modified microbe.

29. The method of Claim 27, wherein said using step comprises using the genetically modified microbe to produce an increased level of beta-glucosidase of at least about 10-fold.

30. The method of Claim 27, wherein said using step comprises using the genetically modified microbe to produce an increased level of beta-glucosidase of at least about 40-fold.

31. The method of Claim 27, wherein said using step comprises using the genetically modified microbe to produce an increased level of beta-glucosidase of at least about 120-fold.




FIG. 1




FIG. 2




FIG. 3

4/5




FIG. 4

5/5




FIG. 5

INTERNATIONAL SEARCH REPORT

national Application No

PCT/CA 99/00197

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N1/15 C12N1/21 C12N15/62 C12N15/63 C12N15/67
C12N9/42 C12P1/02 C12P1/04 C12P1/06 //C12R1:885

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PARADIS, F.W. ET AL.: "Expression and secretion of beta-glucuronidase and pertussis toxin S1 by Streptomyces lividans" APPL. MICROBIOL.BIOTECHNOL., vol. 45, 1996, pages 646-651, XP002109008 see page 648, "Construction of expression/secretion vectors"	1-31
Y	VAN GEMEREN, L.A. ET AL.: "The effect of pre- and pro-sequences and multicopy integration on heterologous expression of the Fusarium solani pisi cutinase gene in Aspergillus awamori" APPL. MICROBIOL.BIOTECHNOLOGY, vol. 45, 1996, pages 755-763, XP002093937 see page 758	1-31

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

13 July 1999

Date of mailing of the international search report

06/08/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alt, G

INTERNATIONAL SEARCH REPORT

national Application No
PCT/CA 99/00197

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 27306 A (ALKO GROUP LTD) 31 July 1997 see page 17, third paragraph ---	1-31
Y	WO 90 15860 A (GENENCOR, INC.) 27 December 1990 page 11, lines 1-6 ---	1-31
A	WO 92 10581 A (GENENCOR INTERNATIONAL) 25 June 1992 cited in the application see the whole document ---	1
A	BARNETT, C.C. ET AL.: "Cloning and amplification of the gene encoding an extracellular beta-glucosidase from Trichoderma reesei: Evidence for improved rates of saccharification of cellulosic substrates" BIO/TECHNOLOGY, vol. 9, 1991, pages 562-567, XP002109025 cited in the application see the whole document ---	1
A	WO 93 12237 A (UNILEVER PLC) 24 June 1993 see page 3; Example 3 ---	1
A	GOSH, A. ET AL.: "Cellulase secretion from a hyper-cellulolytic mutant of Trichoderma Reesei Rut-C30" ARCHIVES OF MICROBIOLOGY., vol. 140, no. 2-3, 1984, pages 126-133, XP002109009 see the whole document ---	1
A	KERÄNEN, S. AND PENTILÄ, M.: "Production of recombinant proteins in the filamentous fungus Trichoderma reesei" CURRENT OPINION IN BIOTECHNOLOGY, vol. 6, 1995, pages 534-537, XP002109010 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/CA 99/00197

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9727306 A	31-07-1997	AU 1547797 A CA 2241968 A EP 0876494 A	20-08-1997 31-07-1997 11-11-1998
WO 9015860 A	27-12-1990	AT 170556 T AU 627334 B AU 5846790 A CA 2034487 A DE 69032616 D DE 69032616 T EP 0429628 A ES 2120947 T JP 4500313 T US 5679543 A	15-09-1998 20-08-1992 08-01-1991 17-12-1990 08-10-1998 11-03-1999 05-06-1991 16-11-1998 23-01-1992 21-10-1997
WO 9210581 A	25-06-1992	CA 2097180 A EP 0562003 A FI 932633 A JP 6503960 T	11-06-1992 29-09-1993 09-06-1993 12-05-1994
WO 9312237 A	24-06-1993	AU 3158393 A CA 2123416 A EP 0672149 A FI 942691 A JP 2866739 B JP 7501453 T US 5705358 A ZA 9209539 A	19-07-1993 24-06-1993 20-09-1995 08-06-1994 08-03-1999 16-02-1995 06-01-1998 09-06-1994

THIS PAGE BLANK (OPTION)